








sigvaris

MEDICAL COMPRESSION PRESCRIPTION PAD

Patient	Date
Diagnosis	Number of refills
	Number of pairs
	No substitution

Physician Name	
Physician Signature 	
License Number	Phone

Compression (mmHg)				
<input type="checkbox"/> 15-20	<input type="checkbox"/> 20-30	<input type="checkbox"/> 30-40	<input type="checkbox"/> 40-50	<input type="checkbox"/> 50-60

<input type="checkbox"/> Closed Toe <input type="checkbox"/> Open Toe <input type="checkbox"/> Patient Choice						
						
<input type="checkbox"/> Calf <input type="checkbox"/> w/Grip-top	<input type="checkbox"/> Thigh	<input type="checkbox"/> Pantyhose	<input type="checkbox"/> L <input type="checkbox"/> R Thigh w/ waist	<input type="checkbox"/> Maternity/ Plus Sizes	<input type="checkbox"/> Wraps <input type="checkbox"/> w/Foot piece	<input type="checkbox"/> Armsleeve <input type="checkbox"/> w/Glove <input type="checkbox"/> w/Gauntlet

Donning Devices:

Please see reverse side for indications

- | | | |
|---|---|---|
| <input type="radio"/> Rubber Gloves | <input type="radio"/> SIMON | <input type="radio"/> Magnide® |
| <input type="radio"/> Latex-free Gloves | <input type="radio"/> Sim-Slide® | <input type="radio"/> SIGVARIS Doff'n Donner |
| <input type="radio"/> Textile Gloves | <input type="radio"/> Cotton Underliner | <input type="radio"/> SIGVARIS Doff'n Donner Cone |

Oona

GRADUATED COMPRESSION CHART

Symptoms/Indications	Compression strength at the ankle (mmHg)
CEAP 0/A: <ul style="list-style-type: none"> Extended standing or sitting (travel) Prophylaxis (pregnancy, risk factors) 	15-20 mmHg
CEAP 1/S: <ul style="list-style-type: none"> Heavy, fatigued, tired legs 	
CEAP 1/S: <ul style="list-style-type: none"> Painful, fatigued and aching legs Spider veins/mild edema (due to pregnancy, age, travelling, etc.) Post sclerotherapy of small veins 	20-30 mmHg
CEAP 2/S: <ul style="list-style-type: none"> Mild to moderate varicose veins Elective surgery (sclerotherapy, phlebectomy, vein stripping, orthopaedics) Orthostatis/Postural hypotension 	
CEAP 3/S: <ul style="list-style-type: none"> Moderate/severe edema (pregnancy, risk factors) Surgery (orthopaedics, post fracture, traumatic edema, sclerotherapy, phlebectomy, vein stripping) 	30-40 mmHg
CEAP 4/S: <ul style="list-style-type: none"> Skin changes without ulceration Superficial Phlebitis (red, painful) Deep Vein Thrombosis/ Post Thrombotic Syndrome 	
CEAP 5/S: <ul style="list-style-type: none"> Skin changes with healed ulceration 	40-50 or 50-60 mmHg
CEAP 6/S: <ul style="list-style-type: none"> Skin changes with active ulceration 	
CEAP 4/S: <ul style="list-style-type: none"> Severe Deep Vein Thrombosis/Post Thrombotic Syndrome 	40-50 or 50-60 mmHg
CEAP 5/S: <ul style="list-style-type: none"> Severe skin changes with healed ulceration (recurrent) 	
CEAP 6/S: <ul style="list-style-type: none"> Severe skin changes with active ulceration 	

CONTRAINDICATIONS	CAUTION
<ul style="list-style-type: none"> Arterial insufficiency, intermittent claudication, ischemia Uncontrolled congestive heart failure Acute dermatitis, weeping dermatosis, cutaneous sepsis 	<ul style="list-style-type: none"> Signs of infection Extensive venous ulceration Skin sensitivities or allergies Neuropathy History of diabetes Confinement to bed or non-ambulatory use unless otherwise prescribed by the physician

No liability accepted for non-observance of contraindications and cautions.